


PROTEOMIC ANALYSIS OF PHOSPHORYLATED NUCLEAR PROTEINS UNDERSCORES NOVEL ROLES FOR RAPID ACTIONS OF RETINOIC ACID IN THE REGULATION OF mRNA SPLICING AND TRANSLATION



Emilio J. Laserna¹, M. Luz Valero², Libia Sanz³, Manuel M. Sánchez del Pino², Juan J. Calvete³ and Domingo Baretino¹
1 Biology of Hormone Action Unit. Instituto de Biomedicina de Valencia (CSIC). Jaime Roig, 11. 46010 Valencia (Spain).
2 Proteomics Unit. Centro de Investigación Príncipe Felipe (CIPF). Av. Autopista del Saler, 16. 46023 Valencia (Spain).
3 Structural Proteinomics Unit. Instituto de Biomedicina de Valencia (CSIC). Jaime Roig, 11. 46010 Valencia (Spain).


Retinoic Acid (RA) signaling is mediated by the Retinoic Acid Receptor (RAR), belonging to the Nuclear Hormone Receptor superfamily. In addition to its classical transcriptional actions, RAR also mediates rapid transcription-independent actions (also called non-genomic), consisting in the activation of signal transduction pathways, as the phosphatidylinositol-3-kinase (PI3K) or the ERK MAP-kinase signaling pathways. RA-induced rapid actions play roles in different physiological contexts. As an effort towards understanding the functions of those rapid actions on signaling elicited by RA, we have identified nuclear proteins whose phosphorylation state is rapidly modified by RA treatment in neuroblastoma cells, using a proteomic approach. Our results show that RA treatment led to changes in the phosphorylation patterns in two families of proteins: (i) those related to chromatin dynamics in relation to transcriptional activation, and (ii) those related to mRNA processing and in particular mRNA splicing. We show that treatment of neuroblastoma cells with RA leads to alteration of the regulation of pre-mRNA splicing and mRNA translation, via the activation of signaling pathways. Thus, our results underscore novel functions for the non-genomic signaling elicited by RAR in the regulation of mRNA processing, as part of a cellular response orchestrated by the nuclear receptor RAR. The results shown here contribute to the idea that transcriptional and transcription-independent actions elicited by nuclear hormone receptors are integrated and converge at multiple levels in the regulation of gene expression.

1. Short RA treatments lead to modifications in the phosphorylation patterns of nuclear proteins



Short RA treatments induce the activation of the PI3K signalling pathway and its downstream components mTOR and p70S6 Kinase (Fig. 1A). Western blot with an antibody against phospho-serine/threonine residues within Akt kinase phosphorylation consensus site (RXRXXS/T), revealed that RA treatment increased rapidly the phosphorylation of several Akt substrate proteins, detectable in whole cell extracts (Fig. 1B), as well as in nuclear extract (Fig. 1C). 2-DE western blots demonstrate that RA treatment leads to an increase in the number and intensity of the spots corresponding to phosphorylated Akt substrates in nuclear extracts from neuroblastoma cells (Fig. 1D).


2. A proteomic approach to the identification of RA-induced phosphorylation changes in nuclear proteins



Spot no.	Name (symbol)
1	Nucleophosmin (NPM)
2, 9	Histone H1.5
3, 4	High Mobility Group Box 1 (HMGB1)
5, 6	Heterogeneous nuclear ribonucleoprotein C1/C2 (hnRNP C1/C2)
7	Heterogeneous nuclear ribonucleoprotein K (hnRNP K)
8	Polyadenylate binding Protein-2 (PABP2)

Nuclear phospho-proteins from cells treated for 30 min with RA or vehicle were prepared by affinity chromatography. The two samples were electrophoresed in 2D-gels and the patterns of spots compared. Differentially phosphorylated proteins were isolated and identified by Mass Spectrometry. RA-induced phosphorylation was validated in 2D-western blot.

3. Analysis of RA-induced differentially phosphorylated proteins through the iTRAQ assay




The iTRAQ method is based in the labeling of peptide fractions with reporter tags of different masses. We run in parallel the affinity-purified nuclear phospho-proteins from cells treated during 0, 15 and 30 min with RA. After denaturation and trypsin digestion, the samples were labeled and mixed. The peptides were separated by multi-dimensional nano-liquid chromatography, and the different fractions analyzed by tandem Mass Spec. The proteins are identified on the basis of peptide fragments, and a relative quantification is obtained from the relative levels of each reporter tag.

Phosphorylation increased by RA	Phosphorylation reduced by RA
Chromatin protein homolog 1 (NPM)	ATP-dependent DNA helicase 2 subunit 1 (T74)
Chromatin protein homolog 3 (BAP1)	Barrier-to-autointegration factor 1 (BAP1)
Chromatin protein homolog 5 (BAP1)	ATP-dependent DNA helicase 2 subunit 2 (T74)
Heterogeneous nuclear ribonucleoprotein A1	Heterogeneous nuclear ribonucleoprotein A1
Heterogeneous nuclear ribonucleoprotein A2	Heterogeneous nuclear ribonucleoprotein A2
Heterogeneous nuclear ribonucleoprotein A3	Heterogeneous nuclear ribonucleoprotein A3
Heterogeneous nuclear ribonucleoprotein A4	Heterogeneous nuclear ribonucleoprotein A4
Heterogeneous nuclear ribonucleoprotein A5	Heterogeneous nuclear ribonucleoprotein A5
Heterogeneous nuclear ribonucleoprotein A6	Heterogeneous nuclear ribonucleoprotein A6
Heterogeneous nuclear ribonucleoprotein A7	Heterogeneous nuclear ribonucleoprotein A7
Heterogeneous nuclear ribonucleoprotein A8	Heterogeneous nuclear ribonucleoprotein A8
Heterogeneous nuclear ribonucleoprotein A9	Heterogeneous nuclear ribonucleoprotein A9
Heterogeneous nuclear ribonucleoprotein A10	Heterogeneous nuclear ribonucleoprotein A10
Heterogeneous nuclear ribonucleoprotein A11	Heterogeneous nuclear ribonucleoprotein A11
Heterogeneous nuclear ribonucleoprotein A12	Heterogeneous nuclear ribonucleoprotein A12
Heterogeneous nuclear ribonucleoprotein A13	Heterogeneous nuclear ribonucleoprotein A13
Heterogeneous nuclear ribonucleoprotein A14	Heterogeneous nuclear ribonucleoprotein A14
Heterogeneous nuclear ribonucleoprotein A15	Heterogeneous nuclear ribonucleoprotein A15
Heterogeneous nuclear ribonucleoprotein A16	Heterogeneous nuclear ribonucleoprotein A16
Heterogeneous nuclear ribonucleoprotein A17	Heterogeneous nuclear ribonucleoprotein A17
Heterogeneous nuclear ribonucleoprotein A18	Heterogeneous nuclear ribonucleoprotein A18
Heterogeneous nuclear ribonucleoprotein A19	Heterogeneous nuclear ribonucleoprotein A19
Heterogeneous nuclear ribonucleoprotein A20	Heterogeneous nuclear ribonucleoprotein A20
Heterogeneous nuclear ribonucleoprotein A21	Heterogeneous nuclear ribonucleoprotein A21
Heterogeneous nuclear ribonucleoprotein A22	Heterogeneous nuclear ribonucleoprotein A22
Heterogeneous nuclear ribonucleoprotein A23	Heterogeneous nuclear ribonucleoprotein A23
Heterogeneous nuclear ribonucleoprotein A24	Heterogeneous nuclear ribonucleoprotein A24
Heterogeneous nuclear ribonucleoprotein A25	Heterogeneous nuclear ribonucleoprotein A25
Heterogeneous nuclear ribonucleoprotein A26	Heterogeneous nuclear ribonucleoprotein A26
Heterogeneous nuclear ribonucleoprotein A27	Heterogeneous nuclear ribonucleoprotein A27
Heterogeneous nuclear ribonucleoprotein A28	Heterogeneous nuclear ribonucleoprotein A28
Heterogeneous nuclear ribonucleoprotein A29	Heterogeneous nuclear ribonucleoprotein A29
Heterogeneous nuclear ribonucleoprotein A30	Heterogeneous nuclear ribonucleoprotein A30
Heterogeneous nuclear ribonucleoprotein A31	Heterogeneous nuclear ribonucleoprotein A31
Heterogeneous nuclear ribonucleoprotein A32	Heterogeneous nuclear ribonucleoprotein A32
Heterogeneous nuclear ribonucleoprotein A33	Heterogeneous nuclear ribonucleoprotein A33
Heterogeneous nuclear ribonucleoprotein A34	Heterogeneous nuclear ribonucleoprotein A34
Heterogeneous nuclear ribonucleoprotein A35	Heterogeneous nuclear ribonucleoprotein A35
Heterogeneous nuclear ribonucleoprotein A36	Heterogeneous nuclear ribonucleoprotein A36
Heterogeneous nuclear ribonucleoprotein A37	Heterogeneous nuclear ribonucleoprotein A37
Heterogeneous nuclear ribonucleoprotein A38	Heterogeneous nuclear ribonucleoprotein A38
Heterogeneous nuclear ribonucleoprotein A39	Heterogeneous nuclear ribonucleoprotein A39
Heterogeneous nuclear ribonucleoprotein A40	Heterogeneous nuclear ribonucleoprotein A40
Heterogeneous nuclear ribonucleoprotein A41	Heterogeneous nuclear ribonucleoprotein A41
Heterogeneous nuclear ribonucleoprotein A42	Heterogeneous nuclear ribonucleoprotein A42
Heterogeneous nuclear ribonucleoprotein A43	Heterogeneous nuclear ribonucleoprotein A43
Heterogeneous nuclear ribonucleoprotein A44	Heterogeneous nuclear ribonucleoprotein A44
Heterogeneous nuclear ribonucleoprotein A45	Heterogeneous nuclear ribonucleoprotein A45
Heterogeneous nuclear ribonucleoprotein A46	Heterogeneous nuclear ribonucleoprotein A46
Heterogeneous nuclear ribonucleoprotein A47	Heterogeneous nuclear ribonucleoprotein A47
Heterogeneous nuclear ribonucleoprotein A48	Heterogeneous nuclear ribonucleoprotein A48
Heterogeneous nuclear ribonucleoprotein A49	Heterogeneous nuclear ribonucleoprotein A49
Heterogeneous nuclear ribonucleoprotein A50	Heterogeneous nuclear ribonucleoprotein A50
Heterogeneous nuclear ribonucleoprotein A51	Heterogeneous nuclear ribonucleoprotein A51
Heterogeneous nuclear ribonucleoprotein A52	Heterogeneous nuclear ribonucleoprotein A52
Heterogeneous nuclear ribonucleoprotein A53	Heterogeneous nuclear ribonucleoprotein A53
Heterogeneous nuclear ribonucleoprotein A54	Heterogeneous nuclear ribonucleoprotein A54
Heterogeneous nuclear ribonucleoprotein A55	Heterogeneous nuclear ribonucleoprotein A55
Heterogeneous nuclear ribonucleoprotein A56	Heterogeneous nuclear ribonucleoprotein A56
Heterogeneous nuclear ribonucleoprotein A57	Heterogeneous nuclear ribonucleoprotein A57
Heterogeneous nuclear ribonucleoprotein A58	Heterogeneous nuclear ribonucleoprotein A58
Heterogeneous nuclear ribonucleoprotein A59	Heterogeneous nuclear ribonucleoprotein A59
Heterogeneous nuclear ribonucleoprotein A60	Heterogeneous nuclear ribonucleoprotein A60
Heterogeneous nuclear ribonucleoprotein A61	Heterogeneous nuclear ribonucleoprotein A61
Heterogeneous nuclear ribonucleoprotein A62	Heterogeneous nuclear ribonucleoprotein A62
Heterogeneous nuclear ribonucleoprotein A63	Heterogeneous nuclear ribonucleoprotein A63
Heterogeneous nuclear ribonucleoprotein A64	Heterogeneous nuclear ribonucleoprotein A64
Heterogeneous nuclear ribonucleoprotein A65	Heterogeneous nuclear ribonucleoprotein A65
Heterogeneous nuclear ribonucleoprotein A66	Heterogeneous nuclear ribonucleoprotein A66
Heterogeneous nuclear ribonucleoprotein A67	Heterogeneous nuclear ribonucleoprotein A67
Heterogeneous nuclear ribonucleoprotein A68	Heterogeneous nuclear ribonucleoprotein A68
Heterogeneous nuclear ribonucleoprotein A69	Heterogeneous nuclear ribonucleoprotein A69
Heterogeneous nuclear ribonucleoprotein A70	Heterogeneous nuclear ribonucleoprotein A70
Heterogeneous nuclear ribonucleoprotein A71	Heterogeneous nuclear ribonucleoprotein A71
Heterogeneous nuclear ribonucleoprotein A72	Heterogeneous nuclear ribonucleoprotein A72
Heterogeneous nuclear ribonucleoprotein A73	Heterogeneous nuclear ribonucleoprotein A73
Heterogeneous nuclear ribonucleoprotein A74	Heterogeneous nuclear ribonucleoprotein A74
Heterogeneous nuclear ribonucleoprotein A75	Heterogeneous nuclear ribonucleoprotein A75
Heterogeneous nuclear ribonucleoprotein A76	Heterogeneous nuclear ribonucleoprotein A76
Heterogeneous nuclear ribonucleoprotein A77	Heterogeneous nuclear ribonucleoprotein A77
Heterogeneous nuclear ribonucleoprotein A78	Heterogeneous nuclear ribonucleoprotein A78
Heterogeneous nuclear ribonucleoprotein A79	Heterogeneous nuclear ribonucleoprotein A79
Heterogeneous nuclear ribonucleoprotein A80	Heterogeneous nuclear ribonucleoprotein A80
Heterogeneous nuclear ribonucleoprotein A81	Heterogeneous nuclear ribonucleoprotein A81
Heterogeneous nuclear ribonucleoprotein A82	Heterogeneous nuclear ribonucleoprotein A82
Heterogeneous nuclear ribonucleoprotein A83	Heterogeneous nuclear ribonucleoprotein A83
Heterogeneous nuclear ribonucleoprotein A84	Heterogeneous nuclear ribonucleoprotein A84
Heterogeneous nuclear ribonucleoprotein A85	Heterogeneous nuclear ribonucleoprotein A85
Heterogeneous nuclear ribonucleoprotein A86	Heterogeneous nuclear ribonucleoprotein A86
Heterogeneous nuclear ribonucleoprotein A87	Heterogeneous nuclear ribonucleoprotein A87
Heterogeneous nuclear ribonucleoprotein A88	Heterogeneous nuclear ribonucleoprotein A88
Heterogeneous nuclear ribonucleoprotein A89	Heterogeneous nuclear ribonucleoprotein A89
Heterogeneous nuclear ribonucleoprotein A90	Heterogeneous nuclear ribonucleoprotein A90
Heterogeneous nuclear ribonucleoprotein A91	Heterogeneous nuclear ribonucleoprotein A91
Heterogeneous nuclear ribonucleoprotein A92	Heterogeneous nuclear ribonucleoprotein A92
Heterogeneous nuclear ribonucleoprotein A93	Heterogeneous nuclear ribonucleoprotein A93
Heterogeneous nuclear ribonucleoprotein A94	Heterogeneous nuclear ribonucleoprotein A94
Heterogeneous nuclear ribonucleoprotein A95	Heterogeneous nuclear ribonucleoprotein A95
Heterogeneous nuclear ribonucleoprotein A96	Heterogeneous nuclear ribonucleoprotein A96
Heterogeneous nuclear ribonucleoprotein A97	Heterogeneous nuclear ribonucleoprotein A97
Heterogeneous nuclear ribonucleoprotein A98	Heterogeneous nuclear ribonucleoprotein A98
Heterogeneous nuclear ribonucleoprotein A99	Heterogeneous nuclear ribonucleoprotein A99
Heterogeneous nuclear ribonucleoprotein A100	Heterogeneous nuclear ribonucleoprotein A100

The iTRAQ assay resulted in the identification of 63 proteins whose phosphorylation changed as effect of RA treatment.

4. iTRAQ: GeneOntology analysis of RA-induced differentially phosphorylated proteins




Among the proteins whose phosphorylation is modified by RA, the proteins involved in RNA metabolism are majority. The percentage of proteins involved in mRNA splicing is remarkable. Therefore we have postulated the following

HYPOTHESIS:


RA, through the rapid activation of signaling pathways could regulate mRNA processing/splicing as part of a cellular response orchestrated by the nuclear receptor RAR

5. RA influences the regulation of alternative splicing via activation of signaling pathways




Using a splicing reporter minigene based on E1A (Fig. 5A) we could show that RA treatment alters the pattern of alternative splicing. This regulation is sensitive to PI3K inhibitors, but not to MAP kinase inhibitors. (Figs. 5B and C). The interaction between the SR protein SF2/ASF and the U1-70K spliceosome protein is increased by RA treatment, as shown in immunoprecipitation experiments (Fig. 5D).

6. RA regulates translation through the activation of signaling pathways



By using a translation reporter gene containing a binding site for SF2/ASF (Fig. 6A) we show RA-induced increase in translation, that was sensitive to inhibitors of both PI3K and ERK-MAP kinase pathways (Fig. 6B). RA increases the phosphorylation of 4E-BP1, an inhibitor of cap-binding eIF4E, through the activation of PI3K (Fig. 6C). However additional mechanisms must exist to explain the effects of MEK inhibitor U0126, that does not interfere the activation of PI3K pathway (Fig. 6D).

7. RA treatment modifies the alternative splicing pattern on endogenous genes in neuroblastoma cells



Preliminary data obtained with Affymetrix Human Gene 1.0ST microarray, which uses exon-specific probes, indicate that RA-induced changes in the alternative splicing pattern take place in a significant number of genes. The figure shows examples of RA-induced exon-inclusion as well as a complex pattern including both exon inclusion and skipping induced by RA

8. Conclusions

We provide the first evidence that RA, through the activation of signalling pathways, could regulate mRNA splicing and translation. The results shown here contribute to the idea that transcriptional and transcription-independent actions elicited by nuclear hormone receptors are integrated and converge at multiple levels in the regulation of gene expression.